

兔抗 PLEKHG6 多克隆抗体

- 中文名称: 兔抗 PLEKHG6 多克隆抗体
- 英文名称: Anti-PLEKHG6 rabbit polyclonal antibody
- 别名: MyoGEF
- 相关类别: 一抗
- 储存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 抗原: PLEKHG6
- 反应种属: Human, Mouse
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	PLEKHG6 functions as a guanine nucleotide exchange fact or activating the small GTPase RHOA, which, in turn, indu ces myosin filament formation. It activates RHOG but not activate RAC1, or to a much lower extent than RHOA an d RHOG. Part of a functional unit, involving MyoGEF, MY H10 and RHOA, at the cleavage furrow to advance furro w ingression during cytokinesis. In epithelial cells, require d for the formation of microvilli and membrane ruffles o n the apical pole. Along with EZR, required for normal m acropinocytosis.
Applications:	ELISA, WB, IHC
Name of antibody:	PLEKHG6
Immunogen:	Fusion protein of human PLEKHG6



www.shjning.com

Full name:	pleckstrin homology domain containing, family G (with R hoGef domain) member 6
Synonyms :	MyoGEF
SwissProt:	Q3KR16
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human lung cancer and human cervical cancer
IHC Recommend dilution:	100-300
WB Predicted band size:	89 kDa
WB Positive control:	231 cells
WB Recommended dilution:	500-2000





